

Natural Resource Management



Natural Resource Management

Fire Management.

Background.

Wildfires have been a component of upland ecosystems on Cape Cod for thousands of years. Prior to European settlement, Cape vegetation was frequently burned by both natural and anthropogenic wildfires (Backman, 1984; Day, 1953; Patterson and Sassman, 1988), and it is likely that fire has played a role in the spatial heterogeneity of upland vegetative cover on the outer Cape. Since the establishment of Cape Cod National Seashore in 1961, however, CACO staff have aggressively suppressed all wildfires within the seashore's boundary, limiting total burned acres to less than 40 hectares in as many years. The effect of this full-suppression policy on forest fuels and vegetative communities within the seashore remains unknown, but it is hypothesized that complete fire suppression may lead to more uniform forest cover, the decrease and eventual disappearance of fire-dependent communities such as grasslands and heathlands, an increase in surface water acidity and an unprecedented accumulation of organic matter with the potential to fuel an uncontrollable and highly destructive wildfire.

An average of 10 to 15 ignitions occur each year at CACO, all the result of human carelessness or mischief (Dosmann and Patterson, 1990). When paired with climate and forest fuel factors, this human impact can create the potential for hazardous wildfires on the outer Cape. The abundance of black huckleberry (*Gaylussacia baccata*) in much of the park's pine-oak forests also presents a fire management concern, because it contains volatile oils that burn explosively when ignited. Since 1985, CACO and cooperative research units at the University of Massachusetts have been conducting prescribed burning and collections of vegetation response data in a 7-hectare site in South Truro. This research has, in turn, led to the development of several Cape-specific fuel models, which are currently being tested under prescribed burning conditions on one- to five-acre plots at the Truro site. Burns took place in 1999, 2000 and 2001, and are planned to continue through 2004.

Research Needs.

Map Hazard Fuels: The accumulation of forest fuels at CACO has increased dramatically in recent years, largely as a result of insect, wind and disease damage. Pitch pine (*Pinus rigida*) communities have been particularly hard hit, with turpentine beetles (*Dendroctonus* sp.) leading to blue stain fungal infection in stands as large as twenty hectares. Updated hazard fuel mapping and sampling, including biomass measurements, are needed in order to make informed decisions regarding fire management in the park.

Assess Forest Pest Damage: The United States Forest Service (USFS) coordinates annual nationwide gypsy moth monitoring using a variety of sampling techniques, including traps, burlap bands and aerial surveys. Gypsy moth populations appear to be on the rise in Massachusetts, and continued monitoring using the USFS methods is necessary in order to track changes in the local abundance of this devastating invasive species and to formulate park-specific gypsy moth management actions. Annual surveys of browntail

Natural Resource Management ---

Fire Management, continued.

moths and other forest pests are also needed in order to inventory and assess the damage caused by insect pests in CACO forests. When possible, survey results should be linked to environmental factors such as temperature and drought status in order to provide resource managers with a tool for predicting damaging infestations and subsequent hazard fuel accumulations.

(See related project description under “The Browntail Moth” and “The Gypsy Moth” in the Wildlife Ecology chapter.)

Investigate Suppression Impacts: Although it is hypothesized that CACO’s policy of complete wildfire suppression could precipitate a decline in habitat heterogeneity or result in a major uncontrolled wildfire with extensive natural and cultural repercussions, little quantitative information about the impacts of fire suppression on the outer Cape actually exists. A thorough investigation of fire suppression impacts on Cape ecosystems is needed, followed by long term monitoring of CACO’s plant communities and computer modeling of future vegetation trends. The fuel loading transects completed by Patterson, et al. in 1984 should be re-surveyed, and research should be undertaken to determine the present level of fuel loading in the park, as well as the average rate of accumulation.

Test Fuel Models for Coastal Pine Communities: Five years of research on huckleberry (Patterson, et al., 1984) has produced data on vegetation response to fire, fire rate-of-spread and fireline intensity for prescribed burns in the huckleberry understory of oak-pine forest, and ongoing research on the effects of fire on coastal pine communities has resulted in the development of CACO-specific fuel models with huckleberry components. Field validation trials are currently in progress at one site in South Truro; however, at least four discrete areas need to be tested in order to check the validity of these models. Continued testing of custom fuel models is thus necessary.

Research Cited.

Backman, A.E. 1984. 1000-year record of fire-vegetation interaction in the northeastern United States: A comparison between coastal and inland regions. Master’s Thesis. University of Massachusetts, Amherst, MA.

Day, G. 1953. The indian as an ecological factor in the northeastern forests. *Ecology* 34:329-346.

Dosmann, M. and W.A. Patterson III. 1990. The woods roads of Cape Cod National Seashore: Access and management for emergency vehicle use. Department of Forestry and Wildlife Management, University of Massachusetts, Amherst, MA.

Natural Resource Management

Fire Management, continued.

Patterson, W.A. and K.E. Sassman. 1988. Indian fires in the prehistory of New England, in *Holocene Human Ecology in Northeastern North America* (G.P. Nicholas ed.) Plenum Publishing Corporation, New York, NY.

Patterson, W.A., K.E. Saunders and L.J. Horton. 1984. Fire regimes of Cape Cod National Seashore. USDI – NPS Office of Scientific Studies Report OSS 83-1.

--- *Natural Resource Management*

Land Use Mapping.

Background.



The last thirty years have brought unprecedented population growth to Cape Cod, along with a boom of residential and commercial development that shows little sign of slowing. Between 1970 and 2000, the number of year-round Cape Cod residents doubled to 200,000, the number of Cape housing units increased by a dramatic 76 %, and despite the ability of Cape Cod National Seashore to protect pristine land inside its

boundaries from major development, increasing numbers of privately-owned seasonal cottages within the park have been, and continue to be, redeveloped and converted to year-round residences. Forest clearance, extended pavement and new roads, additional on-site wastewater disposal and increased use of pesticides and fertilizers in areas within and adjacent to CACO all pose a serious threat to the outer Cape's aquatic and terrestrial resources. Rather than attempting to assess each development project individually, Geographic Information System mapping and analysis are expected to provide a more systematic, comprehensive and scientifically credible way to assess the cumulative impacts of development on outer Cape Cod.

Research Needs.

Regular land use mapping and analysis, using CACO's GIS, is needed to monitor and assess rapid land use changes on the outer Cape. Although preliminary data can be acquired from MassGIS (Commonwealth of Massachusetts, EOE -- land use, wetlands and conservation lands data from 1999 are now available for internet download), sources of impact such as new construction, subdivisions, large paved areas, lawns, underground storage tanks and new septic systems need to be mapped from aerials, field visits and assessor's records. The GIS analysis should include hydrologic (groundwater and surface water flow direction and velocities) and geologic (soils and surficial geology) information to predict development effects on downgradient wetlands, ponds and estuaries. Additionally, specific GIS Software tools and applications for predicting groundwater effects need to be identified and tested for use at CACO.

Natural Resource Management

Long-Term Weather Database.



Background.

The success of nearly all ecological field research and monitoring at Cape Cod National Seashore, as well as the investigation of oil spills, wildfires, insect dispersion and other unexpected impacts to CACO natural resources, is dependent upon accurate and accessible meteorological information. Presently, precipitation

data are recorded continuously at the Truro National Atmospheric Deposition site and Race Point Ranger Station, and National Oceanic and Atmospheric Administration observations are collected at weather stations in Provincetown and Chatham. Fire weather observations including temperature, RH, precipitation, fuel stick moisture, and wind speed and direction are collected hourly during fire season on a data logger, which is downloaded as needed for fire indices determinations.

Research Needs.

A standardized protocol for monitoring meteorological and atmospheric conditions at CACO is currently being developed in partnership with the United States Geological Survey. Upon its completion, long-term monitoring needs to be initiated and subsequent data integrated, along with past and present fire weather observations, into a comprehensive weather database. As an important tool for enhancing research and information across the resource management spectrum, the database should be formatted for easy retrievability by CACO resource managers.

Natural Resource Management

Vegetation Mapping.

Background.

Up-to-date vegetation maps are an important component of vegetation and fuels management, and are vital to predicting ecological change stemming from natural processes, unplanned human intervention and habitat restoration projects. Accurate historical vegetation cover maps are also necessary to track changes in plant communities over time and to provide context for other kinds of habitat mapping and research within Cape Cod National Seashore's Inventory & Monitoring program. Vegetation cover mapping needs to be completed at ten-year intervals (and in areas of rapid change, every five years or less) in order to accurately correlate information with other monitoring protocols. With the cooperation of the University of Massachusetts, an updated vegetation map is currently being generated based on aerial photos from 2000; the next update should occur by 2010. Traditional hard-copy photogrammetry methods are still being utilized, but as high resolution satellite imagery becomes more available and new remote sensing classification methods build proven track records, these methods may also be considered.

Research Needs.

Restore Historic Maps: Historical cover and vegetation maps are a valuable source of information which, if their methods and accuracy can be verified, may add decades to the long-term record of vegetation and land use changes on outer Cape Cod. Historical land cover and vegetation map sources for the outer Cape exist in various forms: hard-copy vegetation polygon maps from 1958 and 1977 (with partial digital versions), U.S. Coastal Survey Maps from 1836-1868 and 1943, and complete aerial photo sets from 1938, 1947, 1960, 1977 and 1987. Other recent photo sets are also becoming available from the Commonwealth of Massachusetts. Information from each of these sources needs to be digitized, interpreted, checked against contemporary evidence and verified, as well as metadata prepared. The final geographic data standard dictates ArcInfo format, UTM coordinate system, North American Datum 1983 (usable in Arcview 3.2a and subsequent versions of ESRI software).

Verify Current Maps: Current and future vegetation mapping efforts need field verification, and statistical field sampling (including ordination analysis) is needed for a more precise record of vegetation type composition. Once sampling methods have been standardized, field sampling and verification efforts should be repeated at regular intervals. Current data should also be compared to historical information using descriptive methods. Additional query/analysis may be possible based on specific questions of habitat change and wildlife dynamics, such as trends in heathland, wetland and forest composition, threatened species productivity, predator interactions and forest pests.

Natural Resource Management

Additional Projects in Natural Resource Management.

Many natural resource management projects at Cape Cod National Seashore also fall under other areas of study. See below for additional natural resource management projects listed in this catalog under their primary field:

Aquatic Ecology

Aquaculture Impacts on Estuarine Ecosystems, p. 1-1

Estuarine Habitat Restoration, p. 1-7

Groundwater Withdrawal, p. 1-15

Gull Pond Sluiceway, p. 1-17

Invasive Aquatic Species, p. 1-21

Kettle Ponds, p. 1-25

- Survey Invasive Species and Develop an Emergency Response Plan

- Develop Individual Management Plans for Each Pond

- Develop a Comprehensive Kettle Pond Management Plan

- Develop Revegetation Plans

- Study Public Use

Landfill Impacts on Groundwater, p. 1-37

- Literature Review of Capping Methods

Larvicide Impacts on Native Invertebrates, p. 1-39

Marine Debris Monitoring, p. 1-41

Mercury Contamination of Aquatic Environs, p. 1-43

Atmospheric Monitoring

Air Quality Monitoring, p. 2-1

Coastal Geomorphology

Physical Oceanographic Processes, p. 3-1

Plant Ecology

Heathlands, p. 4-1

- Test Management Techniques

- Develop Management Plan

Landscape Revegetation, p. 4-5

- Evaluate Dune Planting Program

Non-Native Plant Species, p. 4-7

- Develop Management Plan

Non-Vascular Plant Inventory, p. 4-9

- Assess Mushroom Harvest

Natural Resource Management —————

Additional Projects in Natural Resource Management, continued.

Wildlife Ecology

The Browntail Moth, p. 5-1

Study Browntail Impacts on Native Vegetation

The Gypsy Moth, p. 5-7

Hunting Impacts, p. 5-9

Off-Road Vehicle Traffic Impacts on Invertebrates, p. 5-15

Marsh-Dwelling Shorebirds, p. 5-11

Evaluate the Impacts of Aquaculture on Fisheries and Shorebird Habitat Rare

Invertebrates, p. 5-17

Evaluate Northeastern Beach Tiger Beetle Reintroduction Potential

Cultural Resource Management

Cranberry Bog Restoration, p. 7-3

Natural Resource Management

Selected Bibliography for Natural Resource Management.

Adamas, P. and K. Brandt. 1998. Update of impacts on quality of inland wetlands of the United States: A survey of indicators, techniques and applications of community level biomonitoring data. EPA/600/3-90/073. U.S. Environmental Protection Agency, Washington, D.C.

Anderson. 1989. Determination of the efficacy of clipping browntail moth hibernacles as a means of population control. Unpublished report, Cape Cod National Seashore. South Wellfleet, MA.

Appleton, E. 1994. The war on waste: National parks as dumping grounds for hazardous wastes. *National Parks*. 68(9-10): 37.

Association for the Preservation of Cape Cod, Inc. 1990. Cape Cod Critical Habitat Atlas. Orleans, MA.

----. 1991. More Than Sandy Beaches: A Guide to the Rare and Threatened Habitats of Cape Cod. Orleans, MA.

Atlantic Coast Piping Plover Recovery Team. 1995. Revised Recovery Plan, Piping Plover (*Charadrius melodus*), Atlantic Coast Population. United States Fish and Wildlife Service. Technical/Agency Draft. Region #5, Hadley, MA.

Backman, A.E. 1984. 1000-year record of fire-vegetation interaction in the northeastern United States: A comparison between coastal and inland regions. Master's Thesis. University of Massachusetts, Amherst, MA.

Barlow, P.M. 1994. Particle tracking analysis of contributing areas of public supply wells in simple and complex flow systems, Cape Cod, Massachusetts. USGS Open File Report 93-159. U.S. Geological Survey, Reston, VA.

Beatty, L.L., B.L. Nowicki, A. Keller, R.A. Wahle, C.L. LaBash, and P. V. August. 1994. A Plan for Inventory and Monitoring of Estuarine Resources at Cape Cod National Seashore. NPS Technical Report. NPS/NAROSS/NRTR-94-21. NPS North Atlantic Region, Boston, MA.

Brandes, M. 1977. Effective phosphorus removal by adding alum to septic tank. *Jour. Water Pollution Control Fed.*, 49(11): 2285-2296.

Brownlow, A.H. 1979. Cape Cod environmental atlas. Department of Geology, Boston University, Boston, MA.

Natural Resource Management —————

Selected Bibliography for Natural Resource Management, continued.

Cambareri, T.C. 1986. Hydrogeology and hydrochemistry of a sewage effluent plume in the Barnstable Outwash of the Cape Cod Aquifer, Hyannis, Massachusetts. Master's Thesis. University of Massachusetts, Amherst, MA.

Cambareri, T., G. Belfit, D. Janik, P. Irvin, B. Campbell, D. McCaffery and A. Carbonell. 1989a. Truro/Provincetown aquifer assessment and groundwater protection plan. Cape Cod Planning and Economic Development Commission Water Resources Office, Barnstable, MA.

Cambareri, T., G. Belfit and D. Janik. 1989. Hydrogeologic assessment of the Truro Landfill, Truro, Massachusetts. Cape Cod Planning and Economic Development Commission Water Resources Office, Barnstable, MA.

Cantor, L.W. and R.C. Knox. 1985. Septic tank system effects on groundwater quality. Lewis Publishers, Chelsea, MI.

Cape Cod Commission. 1991. Cape Cod Commission Regional policy plan and atlas. Barnstable, MA.

----. 1997. Final Report of the Lower Cape Water Management Task Force. Barnstable, MA.

Cape Cod Commission and Land Use Collaborative. 1997. The Outer Cape Capacity Study. 4 vols. Barnstable, MA.

Cape Cod Planning and Economic Development Commission. 1978. Environmental Impact Statement and 208 Water quality management plan for Cape Cod, vol. 1 and 2.

----. 1989. Truro/Provincetown Aquifer Assessment and Groundwater Protection Plan. Water Resources Office, Barnstable, MA.

Coastal Engineering Co., Inc. 1992. Initial site assessment report (ISA) of Orleans Sanitary Landfill. Orleans, MA.

----. 1993. Initial site assessment report (ISA) of Wellfleet Sanitary Landfill. Wellfleet, MA.

Colman, J.A., P.K. Weiskel and J.W. Portnoy. 2000. Ground-water nutrient transport to estuaries and freshwater ponds, Cape Cod National Seashore, Massachusetts: A flow path and geochemical simulation approach. Proposal for funding in 2001 under the National Park Service/USGS Water Resources Division Water-quality Monitoring and Assessment (WAQAM) Partnership.

Natural Resource Management

Selected Bibliography for Natural Resource Management, continued.

Curley, J.R., R.P. Lawton, D.K. Whitaker, and J.M. Hickey. 1972. A study of the marine resources of Wellfleet Harbor. Massachusetts Division of Marine Fisheries. 1972 Apr; Monograph Series, No. 12.

Crary, D.W. Jr. 1987. The effect of prescribed burning and mowing on fuels in *Gaylussacia baccata* communities. Unpublished. Master's Special Project, Department of Forestry and Wildlife Management, University of Massachusetts, Amherst, MA.

Day, G. 1953. The indian as an ecological factor in the northeastern forests. *Ecology* 34:329-346.

Desimone, L.A. and B.L. Howes. 1995. Hydrogeologic, water quality and biogeochemical data collected at a septage treatment facility, Orleans, Cape Cod, Massachusetts, October 1988 through December 1992. USGS Open File Report 95-439, U.S. Geological Survey, Marlborough, MA.

----. 1996. Denitrification and nitrogen transport in a coastal aquifer receiving wastewater discharge. *Environmental Science and Technology*, 30(4).

Dosmann, M. and W.A. Patterson III. 1990. The woods roads of Cape Cod National Seashore: Access and management for emergency vehicle use. Department of Forestry and Wildlife Management, University of Massachusetts, Amherst, MA.

Dunwiddie, P.W. and M.B Adams. 1995. Fire suppression and landscape change on outer Cape Cod. NPS Technical Report. NPS/NESO-RNR/NRTR/96-08.

East Cape Engineering. 1996. Truro landfill initial site assessment report (ISA). East Cape Engineering, Orleans, MA.

Eichner, E.M. and T.C. Cambareri. 1991. Nitrogen loading. Cape Cod Planning and Economic Development Commission Water Resources Office. Technical Report 91-001. Barnstable, MA.

Eichner, E.M., T.C. Cambareri, and K. Livingston. 1996. Groundwater impacts associated with the removal of the tide gate and dike structures on the Pamet River, Truro, Massachusetts. Cape Cod Commission. Barnstable, MA.

Frimpter, M.H. and G.C. Belfit. 1992. Estimation of high ground water levels for construction and land use planning: A Cape Cod, Massachusetts example. Cape Cod Commission Technical Bulletin 92-001. Cape Cod Commission, Barnstable, MA.

Frohlich, R. K. 1991. Geoelectrical survey for locating groundwater pollution from a landfill northwest of Provincetown, Cape Cod, Massachusetts. Department of Geology, University of Rhode Island, Kingston, RI.

Natural Resource Management ---

Selected Bibliography for Natural Resource Management, continued.

Frohlich, R.K. and Urish, D.W. 1994. Use of geoelectrical methods in groundwater pollution surveys in a coastal environment. *J. Appl. Geophysics*. 32:139-154.

Godfrey, P.J., M.A. Benedict and S. Swartz. 1976. Natural resource management plan for Cape Cod National Seashore. NPS-CPSU, University of Massachusetts Report Number 22.

Godfrey, P.J. and L. Stack. In press. Dune revegetation monitoring/Cape Cod National Seashore, Summer 1987. University of Massachusetts, Amherst, MA.

Godfrey, P.J., M. Mattson, M. Walk, P. Kerr, T. Zajicek and A. Ruby III. 1996. The Massachusetts Acid Rain Monitoring Project: Ten years of monitoring Massachusetts lakes and streams with volunteers. Water Resources Research Center. Publication 171. University of Massachusetts, Amherst, MA.

Goetz, W.J. C.R. Harper, C.E. Willis and J.T. Finn. 1991. Effects of land uses and hydrologic characteristics on nitrate and sodium in groundwater. Water Resources Research Center, University of Massachusetts, Amherst, MA.

Godfrey, P.J., K. Galluzzo, N. Price and J.W. Portnoy. 1999. Water Resources Management Plan for Cape Cod National Seashore. National Park Service.

Harris, S.L. and P.A. Steeves. 1994. Identification of potential public water supply areas of the Cape Cod aquifer, Massachusetts, using a geographic information system. USGS Water Resources Investigations Report 94-4156. U.S. Geological Survey, Marlborough, MA.

Killian, J.C. 1985. Cape Cod National Seashore land protection plan. Cape Cod National Seashore, South Wellfleet, MA.

Koske, R.E. and J.W. Gemma. 1992. Restoration of early and late successional dune communities of the Province Lands. Cape Cod National Seashore Technical Report, NPS Coastal Research Center, URI, NPS/NARURI/NRTR-92/03.

Haines, T. 1996. Evaluate mercury contamination in aquatic environments of Acadia National Park and Cape Cod National Seashore, Progress Report. RMP Project Statement ACAD-N-74.000. Acadia National Park, ME.

Harper, C.R. W.J. Goetz and C.E. Willis. 1992. Groundwater protection in mixed land-use aquifers. *Environmental Management*, 16(6): 777-783.

Hershey, A.E., A.R. Lima, G.J. Niemi and R.R. Regal. 1998. Effects of Bti and methoprene on nontarget macroinvertebrates in Minnesota wetlands. *Ecological Applications* 8:41-60.

Natural Resource Management

Selected Bibliography for Natural Resource Management, continued.

Leatherman, S.P. 1981. Dune stabilization of the Provincelands, Cape Cod National Seashore, Massachusetts. Report #52, The Environmental Institute, University of Massachusetts, Amherst, MA.

Leatherman, S.P. and P.J. Godfrey. 1979. The impact of off-road vehicles on coastal ecosystems in Cape Cod National Seashore. Final Report to Cape Cod National Seashore, Contract Number CX-1600-5-0001.

LeBlanc, D.R. 1981. Evaluation of the hydrologic impacts of groundwater withdrawal from a test well site in the Cape Cod National Seashore, Truro, MA. Massachusetts USGS Water Resources Investigations Open File Report.

----. 1984. Sewage plume in a sand and gravel aquifer, Cape Cod, Massachusetts. USGS Water Supply Paper 2218. U.S. Geological Survey, Reston, VA.

Lepore, J.V., C.D. Rugge and K.C. Ahlert. 1990. Provincetown Municipal Landfill Contaminant Assessment. Department of Chemical and Biochemical Engineering, Rutgers University, New Brunswick, NJ.

Manski, D.A. 1990. National park debris monitoring program, Cape Cod National Seashore 1989 Annual Report. Cape Cod National Seashore, South Wellfleet, MA.

----. 1991. National park debris monitoring program, Cape Cod National Seashore 1990 Annual Report. Cape Cod National Seashore, South Wellfleet, MA.

----. 1992. Snow Pond management plan. Cape Cod National Seashore, Wellfleet, MA.

----. 1993. Environmental Assessment for Temporary Use of Beach Access Route at Race Point by Off-Road Vehicles. Cape Cod National Seashore, South Wellfleet, MA.

Martin, L. 1993. Investigations of effects of ground water withdrawals from the Pamet and Chequesset aquifers, Cape Cod National Seashore. Technical Report NPS/NRWRD/NRTR-93/15. National Park Service, Water Resources Division, Fort Collins, CO.

Massachusetts Barrier Beach Task Force. 1994. Guidelines for Barrier Beach Management in Massachusetts.

Massachusetts Department of Environmental Management. 1982. Massachusetts Landscape Inventory: A Survey of the Commonwealth's Scenic Areas. Boston.

Natural Resource Management ---

Selected Bibliography for Natural Resource Management, continued.

Massachusetts Department of Environmental Protection. 1994. Water resources of Cape Cod: Water use, hydrology, and potential changes in ground water levels. Executive Office of Environmental Affairs, Office of Water Resources, Boston, MA.

Massachusetts Department of Environmental Protection. 1997. Truro Landfill summary information. Department of Environmental Protection Landfill Files. Marlborough, MA.

Massachusetts Division of Fisheries and Wildlife. 1990. Abstracts of the 1990 Fish & Wildlife Laws. Massachusetts Division of Fisheries and Wildlife, Boston, MA.

Massachusetts Office of Coastal Zone Management. 1995. The Massachusetts Aquaculture Strategic Plan. Boston.

Masterson, J.P. and P.M. Barlow. 1994. Effects of simulated changes in groundwater pumping and recharge on the hydrology of the sole-source Cape Cod aquifer. U.S. Geological Survey.

----. 2000. Simulated effects of ground-water withdrawals and contaminant migration on the ground-water and surface-water resources of the Cape Cod National Seashore and surrounding areas, Lower Cape Cod, Massachusetts. A proposal to produce an integrated Outer Cape Cod groundwater model. USGS Water Resources Division, Northborough, MA.

Melvin, S.M., L.H. MacIvor, C.R. Griffin, and L.K. Jones. 1992. Piping plover research and management within Cape Cod National Seashore, 1985-91.

Metland Research Group. 1989. Managing adjacent areas: Cape Cod National Seashore case study. Department of Regional Planning/ Landscape Architecture, University of Massachusetts, Amherst, MA.

Mitchell, N.J. and M. Soukup. 1980. Assessment of water resources management alternatives for Cape Cod National Seashore. NPS North Atlantic Region, Office of Scientific Studies, Boston, MA.

----. 1981. Analysis of water resource management: Alternatives with environmental assessment. North Atlantic Region, Office of Scientific Studies, National Park Service, Department of the Interior in Cooperation with Cape Cod National Seashore, Wellfleet, MA.

National Park Service. 1989. Land Protection Plan, Cape Cod National Seashore. Cape Cod National Seashore, South Wellfleet, MA.

Natural Resource Management

Selected Bibliography for Natural Resource Management, continued.

National Park Service. 1990a. Guidelines for natural resources inventory and monitoring. NPS-75, Release No. 1.

----. 1990b. Wildland fire management guidelines. NPS-18.

----. 1994. Fire Management Plan, Cape Cod National Seashore. Cape Cod National Seashore, South Wellfleet, MA.

----. 1996. Cape Cod National Seashore resource management plan. Cape Cod National Seashore, Wellfleet, MA.

----. 1998. Cape Cod National Seashore general management plan. Cape Cod National Seashore, Wellfleet, MA.

Noake, K.D. 1989. Guide to contamination sources for wellhead protection. Cape Cod Aquifer Management Project (CCAMP) Document. US Environmental Protection Agency, Boston, MA.

Noss, R.R. 1989. Recharge area land use and well water quality. The Environmental Institute, Publication 89-2. University of Massachusetts, Amherst, MA.

Noss, R.R., R. Drake and C. Mossman. 1987. Septic tank cleaners: Their effectiveness and impact on groundwater quality. The Environmental Institute, Publication 87-3. University of Massachusetts, Amherst, MA.

Patterson, W.A., K.E. Saunders and L.J. Horton. 1984. Fire regimes of Cape Cod National Seashore. USDI – NPS Office of Scientific Studies Report OSS 83-1.

Persky, J.H. 1986. The relation of ground water quality to housing density, Cape Cod, Massachusetts. USGS Water Resources Investigations Report 86-4093. U.S. Geological Survey, Boston, MA.

Portnoy, J.W. 1984. Salt marsh diking and nuisance mosquito production on Cape Cod, Massachusetts. J. American Mosquito Control Association 44:560-564.

----. 1990a. A prospectus for salt marsh restoration at Hatches Harbor, Provincetown, Massachusetts. Cape Cod National Seashore, Wellfleet, MA.

----. 1990b. Critical elevations for the Hatches Harbor Salt Marsh Restoration Project: A primer. Cape Cod National Seashore, Wellfleet, MA.

----. 1991. Summer oxygen depletion in a diked New England estuary. *Estuaries*; 14(2): 122-129.

Natural Resource Management ---

Selected Bibliography for Natural Resource Management, continued.

Portnoy, J.W. and A.E. Giblin. 1987. Biogeochemical effects of seawater restoration to diked salt marshes. *Ecological Applications*, 7:1054-1063.

Portnoy, J.W., B.A. Nowicki, C.T. Roman, and D. Urish. 1998. The discharge of nitrate-contaminated groundwater from developed shoreline to marsh-fringed estuary. *Water Resources Research*, 34:3095-3104.

Portnoy, J.W., C.T. Roman and M.A. Soukup. 1987. Hydrologic and chemical impacts of diking and draining of a small estuary (Cape Cod National Seashore): Effects on wildlife and fisheries. pp. 253-265 in Whitman, W.R. & W.H. Meredith (eds.). *Proceedings of a Symposium on Waterfowl and Wetlands Management in the Coastal Zone of the Atlantic Flyway*. Delaware Coastal Management Program, Dover, DE.

Portnoy, J.W., C. Phipps and B.A. Samora. 1987. Mitigating the effects of oxygen depletion on Cape Cod National Seashore anadromous fish. *Park Science* 8:12-13.

Portnoy, J.W. and M.A. Soukup. 1988. Can the Cape's diked salt marshes be restored? *The Cape Naturalist*, 17:28-33.

Portnoy, J. and M. Reynolds. 1997. Wellfleet's Herring River: The case for habitat restoration. *Environment Cape Cod*, 1:35-43.

Postma, F.B., A.J. Gold, and G.W. Loomis. 1992. Nutrient and microbial movement from seasonally-used septic systems. *Journal of Environmental Health*, 55(2): 5-10.

Robinson, M.H. 1980. Cape Cod coastal oil spill response plan. Cape Cod Planning and Economic Development Commission. Barnstable, MA.

Roman, C. 1987. An evaluation of alternatives for estuarine restoration management: The Herring River ecosystem (Cape Cod National Seashore). NPS CRU, Rutgers University, New Brunswick, NJ.

Roman, C.T., R.W. Garvine and J.W. Portnoy. 1995. Hydrologic modeling as a predictive basis for ecological restoration of salt marshes. *Environmental Management*, 19(4):559.

Roman, C.T., J. Portnoy, T.C. Cambareri and R. Sobczak. 1996. Potential groundwater withdrawal effects on plant distributions, soils and water chemistry of seasonally-flooded wetlands and kettle ponds of Cape Cod National Seashore. Proposal to the National Park Service, Water Resources Division, Fort Collins, CO.

Samora, B. and M. Whatley. 1986. Resource Management Plan for Cape Cod National Seashore. Cape Cod National Seashore, South Wellfleet, MA.

Natural Resource Management

Selected Bibliography for Natural Resource Management, continued.

Samora, B.A. 1988. Draft management plan for freshwater kettle ponds of Cape Cod National Seashore: Spectacle Pond in Wellfleet and Snow Pond and Round Pond West in Truro. Cape Cod National Seashore, Wellfleet, MA.

SEA Consultants, Inc. 1992. Provincetown sanitary landfill comprehensive site assessment. Cambridge, MA.

----. 1994. Provincetown groundwater report-Revised. Cambridge, MA.

Shortelle, A.B. and E.A. Colburn. 1986. Physical, chemical and biological impacts of liming on a Cape Cod kettle pond. Status Report to the Massachusetts Division of Fisheries and Wildlife, Westborough, MA.

Sinicrope, T.L., P.G. Hine, R.S. Warren and W.A. Niering. 1990. Restoration of an impounded salt marsh in New England. *Estuaries*, 13:25-30.

Smardon, R.C. 1972. Assessing visual-cultural values of inland wetlands in Massachusetts. Master's Thesis. University of Massachusetts, Amherst, MA.

Smardon, R.C. and J.G. Fabos. 1983. A model for assessing visual-cultural values of wetlands: A Massachusetts case study. in R.C. Smardon, ed. *The future of wetlands: Assessing visual-cultural values*. Allanheld, Osmun Publishers, Totowa, NJ.

Sobczak, B. and T. Cambareri. 1995. Lower Cape Water Management Task Force interim report, Fall 1995. Cape Cod Planning Commission, Barnstable, MA.

Soukup, M.A. and J.W. Portnoy. 1986. Impacts from mosquito control-induced sulfur mobilization in a Cape Cod estuary. *Environmental Conservation*, 13(1):47-50.

Town of Eastham. 1996. Eastham local comprehensive plan. Eastham Draft Local Planning Committee and Thomas Planning Consultants, Inc., Boston, MA.

Town of Truro. 1994. Truro comprehensive plan. Truro, MA.

Town of Wellfleet. 1996. Wellfleet local comprehensive plan. Wellfleet, MA.

Town of Provincetown. 1988. Provincetown master plan. Lane Kendig, Inc. Provincetown, MA.

Urish, D.W., M.J. Reilly, R.M. Wright and R.K. Frolich. 1993. Assessment of ground and surface water impacts: Provincetown Landfill and septic disposal site, Provincetown, MA. USDI-NPS Cooperative Park Studies Unit at URO. Technical Report, NPS/NARURI/NRTR-93/01.

Natural Resource Management ---

Selected Bibliography for Natural Resource Management, continued.

U.S. Coast Guard. 1990. Local oil and hazardous substance pollution contingency plan. U.S.D.T.- U.S. Coast Guard Marine Safety Office, Providence, RI.

U.S. Environmental Protection Agency. 1992. Guidance specifying management measures for nonpoint pollution in coastal waters. EPA 840-B-92-002. U.S. Environmental Protection Agency, Washington D.C.

Van Dusen, K. 1988. A Plague of Plastics Threatens the Ocean. *Habitat: Journal of the Maine Audubon Society*. 5(5): 26-8.

Veneman, P.L.M. Unpublished. Onsite sewage treatment and disposal system performance, Environmental and public health aspects.

Walter, D.A., B.A. Rea, K.G. Stollenwerk and J. Savoie. 1996. Geochemical and hydrologic controls on phosphorous transport in a sewage-contaminated sand and gravel aquifer near Ashumet Pond, Cape Cod, Massachusetts. USGS Water Supply Paper 2463. U.S. Geological Survey, Marlborough, MA.

Weintraub, J.M. 1988. Private well contamination in Massachusetts. The Special Legislative Commission on Water Supply. Department of Environmental Protection, Boston, MA.

Weiskel, P.K., B.L. Howes and G.R. Heufelder. 1996. Coliform contamination of a coastal embayment: Sources and transport pathways. *Environmental Science and Technology*, 30: 1872-1881.

Winkler, M.G. 1990. Recent changes in trophic status of Duck and Bennett Ponds relative to possible landfill enrichment. NPS, North Atlantic Regional Office.

Zoto, G. A. and T. Gallagher. 1988. Cape Cod aquifer management project (CCAMP) final report. US Environmental Protection Agency, Boston, MA.